




Eur päisches
Patentamt

Eur pean
Patent Office

Office eur péen
des brevets

EP 99 / 2715



09 / 673836

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-
gen stimmen mit der
ursprünglich eingereichten
Fassung der auf dem näch-
sten Blatt bezeichneten
europäischen Patentanmel-
dung überein.

The attached documents
are exact copies of the
European patent application
described on the following
page, as originally filed.

Les documents fixés à
cette attestation sont
conformes à la version
initialement déposée de
la demande de brevet
européen spécifiée à la
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

98107397.6

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

M.B. RIJLING

DEN HAAG, DEN
THE HAGUE,
LA HAYE, LE

12/04/99



Eur päisches
Patentamt

Eur pean
Patent Office

Office eur péen
d s brevets

Blatt 2 der Besch inigung
Sheet 2 of the certificate
Page 2 de l'attestation

Anmeldung Nr.:
Application no.:
Demande n°: 98107397.6

Anmeldetag:
Date of filing:
Date de dépôt: 23/04/98

Anmelder:
Applicant(s):
Demandeur(s):
Hoechst Marion Roussel Deutschland GmbH
65929 Frankfurt am Main
GERMANY

Bezeichnung der Erfindung:
Title of the invention:
Titre de l'invention:

A process for the conversion of echinocandin class of peptides to their C4-homotyrosine monodeoxy analogues

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s) revendiquée(s)

Staat:
State:
Pays:

Tag:
Date:
Date:

Aktenzeichen:
File no.
Numéro de dépôt:

Internationale Patentklassifikation:
International Patent classification:
Classification internationale des brevets:

C07K7/56


Am Anmeldetag benannte Vertragsstaaten:
Contracting states designated at date of filing: AT/BE/CH/CY/DE/DK/ES/FI/FR/GB/GR/IE/IT/LI/LU/MC/NL/PT/SE
Etats contractants désignés lors du dépôt:

Bemerkungen:
Remarks:
Remarques:

A process for the conversion of echinocandin class of peptides to their C4-homotyrosine monodeoxy analogues

5

This invention relates to a process for the conversion of echinocandin class of peptides of the formula I



(I)


10 wherein W, X, Y, Z, R and R' are as defined herein below :

		<u>W</u>	<u>X</u>	<u>Y</u>	<u>Z</u>	<u>R</u>	<u>R'</u>
1.	Echinocandin B	OH	OH	OH	OH	CH ₃	Linoleoyl
2.	Pneumocandin A ₀	OH	OH	OH	OH	CH ₂ -CO-NH ₂	10,12-Dimethyl- myristoyl
3.	Pneumocandin A ₁	H	OH	OH	OH	CH ₃	"
4.	Pneumocandin A ₂	OH	OH	H	H	CH ₃	"
5.	Pneumocandin B ₀	OH	OH	OH	OH	CH ₃	"
6.	Pneumocandin B ₂	OH	OH	H	H	CH ₃	"
7.	Pneumocandin C ₀	OH	OH	OH	OH	CH ₃	"
8.	Mulundocandin	OH	OH	OH	OH	H	12-Methyl- tetradecanoyl

to their C4-homotyrosine monodeoxy analogues of the formula I, wherein W, X, Y, Z, R and R' are as defined herein below:


5			<u>W</u>	<u>X</u>	<u>Y</u>	<u>Z</u>	<u>R</u>	<u>R'</u>
	1.	Deoxyechinocandin B (Echinocandin C)	OH	H	OH	OH	CH ₃	Linoleoyl
	2.	Deoxypneumocandin A ₀	OH	H	OH	OH	CH ₂ -CO-NH ₂	10,12-Dimethyl- myristoyl
10	3.	Deoxypneumocandin A ₁	H	H	OH	OH	CH ₃	"
	4.	Deoxypneumocandin A ₂	OH	H	H	H	CH ₃	"
	5.	Deoxypneumocandin B ₀	OH	H	OH	OH	CH ₃	"
	6.	Deoxypneumocandin B ₂	OH	H	H	H	CH ₃	"
	7.	Deoxypneumocandin C ₀	OH	H	OH	OH	CH ₃	"
15	8.	Deoxymulundocandin	OH	H	OH	OH	H	12-Methyl tetra- decanoyl,

particularly to a process for the conversion of mulundocandin (compound of the formula II)



(II)

to deoxymulundocandin (compound of the formula III)



(III)

1,3- β -glucan synthesis inhibitors are effective antifungal agents against *Candida albicans* and also *Pneumocystis carini*, an opportunistic organism responsible for an often fatal pneumonitis among HIV patients and other immunocompromised hosts. Of all the structural classes of 1,3- β -glucan synthesis inhibitors, only the echinocandins received considerable attention [Ref : J. Med. Chem. 35, 198-200 (1992)]. Echinocandin class of peptides are cyclic hexapeptides having a lipophilic side chain.


Several methods for the conversion of echinocandins to the corresponding deoxy analogues under acidic conditions have been reported [Ref : Tetrahedron Letts., 33, 4529-4532 (1992); US Patent Appl. No. 222157 dated April 4, 1994]. The above methods involve selective reduction of C4-h₂tyr (homotyrosine) hydroxyl group of echinocandins to their monodeoxy analogues with prior protection / deprotection of the equally facile C5-Orn (ornithine) hydroxyl group.

Mulundocandin [J.Antibiotics, 40, 275-280 and 281-289 (1987)] and deoxymulundocandin [Indian patent No. IN 169830 ; J.Antibiotics, 45, 618-623

(1992)] having antifungal properties were isolated from *Aspergillus sydowii* (Bainier and Sartory) Thom and Church var. Nov. *Mulundensis* Roy (culture no. HIL Y-30462). Deoxymulundocandin was found to possess better antifungal activity than mulundocandin. However, the production of deoxymulundocandin during the fermentation was 200 times less than that of mulundocandin.

We have found out by extensive research and experimentation that echinocandin class of peptides of the formula I may be converted to the corresponding C4-htyr monodeoxy analogues, particularly mulundocandin to deoxymulundocandin under neutral conditions. Accordingly, the object of the present invention is to provide a process for the conversion of echinocandin class of peptides of the formula I to the corresponding C4-homotyrosin monodeoxy analogues, particularly mulundocandin (compound of formula II) to deoxymulundocandin (compound of formula III).

According to the invention, there is provided a process for the conversion of echinocandin class of peptides of the formula I



(I)


wherein W, X, Y, Z, R and R' are as defined herein below :

	<u>W</u>	<u>X</u>	<u>Y</u>	<u>Z</u>	<u>R</u>	<u>R'</u>
1. Echinocandin B	OH	OH	OH	OH	CH ₃	Linoleoyl
2. Pneumocandin A ₀	OH	OH	OH	OH	CH ₂ -CO-NH ₂	10,12-Dimethyl- myristoyl
3. Pneumocandin A ₁	H	OH	OH	OH	CH ₃	"
4. Pneumocandin A ₂	OH	OH	H	H	CH ₃	"
5. Pneumocandin B ₀	OH	OH	OH	OH	CH ₃	"
6. Pneumocandin B ₂	OH	OH	H	H	CH ₃	"
7. Pneumocandin C ₀	OH	OH	OH	OH	CH ₃	"
8. Mulundocandin	OH	OH	OH	OH	H	12-Methyl- tetradecanoyl

to their C4-homotyrosine monodeoxy analogues of the formula I, wherein W, X, Y, Z, R and R' are as defined herein below:


	<u>W</u>	<u>X</u>	<u>Y</u>	<u>Z</u>	<u>R</u>	<u>R'</u>
1. Deoxyechinocandin B (Echinocandin C)	OH	H	OH	OH	CH ₃	Linoleoyl
2. Deoxypneumocandin A ₀	OH	H	OH	OH	CH ₂ -CO-NH ₂	10,12-Dimethyl- myristoyl
3. Deoxypneumocandin A ₁	H	H	OH	OH	CH ₃	"
4. Deoxypneumocandin A ₂	OH	H	H	H	CH ₃	"
5. Deoxypneumocandin B ₀	OH	H	OH	OH	CH ₃	"
6. Deoxypneumocandin B ₂	OH	H	H	H	CH ₃	"
7. Deoxypneumocandin C ₀	OH	H	OH	OH	CH ₃	"
8. Deoxymulundocandin	OH	H	OH	OH	H	12-Methyl tetra- decanoyl

particularly to a process for the conversion of mulundocandin (compound of the formula II



(II)

5 to deoxymulundocandin (compound of the formula III)



which consists of a single step selective reduction of C4-htyr (homotyrosine) hydroxyl group of echinocandins to their monodeoxy analogues particularly under neutral conditions without prior protection / deprotection of the equally facile C5-Orn

(ornithine) hydroxyl group and purification of the monodeoxy compound from the crude reaction mixture.

The conversion of echinocandins to their monodeoxy analogues by selective reduction at C4-htyr may be effected by hydrogenolysis with Raney nickel in solvents such as methanol, ethanol, or dioxane at pH 3-9. Preferably, the selective reduction is carried out by hydrogenolysis with Raney nickel in ethanol at pH 7 and room temperature in the ratio of 6.8 ml Raney nickel per millimole of mulundocandin.

- 10 The monodeoxy compounds of the invention may, for example, be purified from the crude reaction mixture as follows :

By fractionation using normal phase chromatography (using alumina or silica gel as stationary phase and eluents such as petroleum ether, ethyl acetate, dichloromethane, chloroform, methanol or combinations thereof), reverse phase chromatography (using reverse phase silica gel like dimethyloctadecylsilylsilica gel, also called RP-18 or dimethyloctylsilylsilica gel also called RP-8 as stationary phase and eluents such as water, buffers such as phosphate, acetate, citrate (pH 2-8) and organic solvents such as methanol, acetonitrile, acetone, tetrahydrofuran or combination of the solvents), gel permeation chromatography - using resins such as "Sephadex LH-20®" (Pharmacia Chemical Industries, Sweden), TSKgel Toyopearl HW (TosoHaas, Tosoh Corporation, Japan) in solvents such as methanol, chloroform or ethyl acetate or their combination or Sephadex G-10 and G-25 in water; or by counter-current chromatography using a biphasic eluent system made up of two or more solvents such as water, methanol, ethanol, *iso*-propanol, *n*-propanol, tetrahydrofuran, acetone, acetonitrile, methylene chloride, chloroform, ethylacetate, petroleum ether, benzene and toluene. These techniques may be used repeatedly or a combination of the different techniques may be used. Counter-current chromatography (liquid-liquid chromatography) using a biphasic eluent system on ITO coil is preferred for purification of the compounds of the invention.

The following experimental example is illustrative of the present invention but not limitative of the scope thereof.

Example 1

5 Mulundocandin (220 mg, 2.2 mM) in ethanol (8 ml)) was stirred with 15 ml of W-2 Raney nickel (pH 7) in ethanol (30 ml) for 3 hours at room temperature. After standing for 15 minutes the supernatant solution was decanted and Raney nickel washed with 3 x 30 ml. ethanol with stirring and filtered. Combined ethanolic solutions were concentrated by distillation under a reduced pressure of 60-70
10 mm/Hg at 35° C to obtain 160 mg (75%) of crude deoxymulundocandin as a slightly green solid.

The crude product was purified by liquid-liquid chromatography on ITO coil using upper layer of CH₂Cl₂ : MeOH : *n*-PrOH : H₂O as the stationary phase and the lower
15 layer as the mobile phase in an ascending mode. The coils (15 + 25 + 215 ml) were connected in series and a flow rate of 0.6 ml/min. at a piston stroke of 60 and pressure 0.5 bars was maintained. The purification of deoxymulundocandin was monitored both by bioactivity against *Candida albicans* and *Aspergillus niger* and by analytical High Pressure Liquid Chromatography (HPLC) [column : (10 x 0.4 cm + 3
20 x 0.4 cm) ODS-Hypersil, 10μ; mobile phase: 50:50 CH₃CN : H₂O ; flow rate : 1 ml/min; Wavelength : 220 nm.) The fractions (4.5 ml each) containing deoxymulundocandin were combined, concentrated by distillation under a reduced pressure of 60-70 mm/Hg at 35°C and lyophilized to yield pure deoxymulundocandin [65 mg (30% yield)]. Also recovered during the above
25 purification of deoxymulundocandin was unreacted mulundocandin in 10% yield.

The semi-synthetic deoxymulundocandin was identical in all respects to the naturally isolated compound and the physico-chemical data is given in Table 1.


TABLE 1

	Appearance :	White powder
	Melting point :	170-172°C
5	$[\alpha]_D$:	- 36.6° (c 0.25, MeOH)
	HPLC RT :	4.42 min
	FAB-MS (Fast Atom: Bombardment mass)	1014.7 (M + Na) ⁺
	¹ H NMR (300 MHz, : CD ₃ OD)	Figure 1 of the accompanying drawings
10	¹³ C NMR (75 MHz, : CD ₃ OD)	Figure 2 of the accompanying drawings

Claims:

1. A process for the conversion of echinocandin class of peptides of the formula

5 I



(I)

wherein W, X, Y, Z, R and R' are as defined herein below :

10

		<u>W</u>	<u>X</u>	<u>Y</u>	<u>Z</u>	<u>R</u>	<u>R'</u>
	1. Echinocandin B	OH	OH	OH	OH	CH ₃	Linoleoyl
	2. Pneumocandin A ₀	OH	OH	OH	OH	CH ₂ -CO-NH ₂	10,12-Dimethyl-myristoyl
15	3. Pneumocandin A ₁	H	OH	OH	OH	CH ₃	"
	4. Pneumocandin A ₂	OH	OH	H	H	CH ₃	"
	5. Pneumocandin B ₀	OH	OH	OH	OH	CH ₃	"
	6. Pneumocandin B ₂	OH	OH	H	H	CH ₃	"
	7. Pneumocandin C ₀	OH	OH	OH	OH	CH ₃	"
20	8. Mulundocandin	OH	OH	OH	OH	H	12-Methyl-tetradecanoyl

to their C4-homotyrosine monodeoxy analogues of the formula I, wherein W, X, Y, Z, R and R' are as defined herein below

		<u>W</u>	<u>X</u>	<u>Y</u>	<u>Z</u>	<u>R</u>	<u>R'</u>
5	1. Deoxyechinocandin B (Echinocandin C)	OH	H	OH	OH	CH ₃	Linoleoyl
	2. Deoxypneumocandin A ₀	OH	H	OH	OH	CH ₂ -CO-NH ₂	10,12-Dimethyl- myristoyl
	3. Deoxypneumocandin A ₁	H	H	OH	OH	CH ₃	"
10	4. Deoxypneumocandin A ₂	OH	H	H	H	CH ₃	"
	5. Deoxypneumocandin B ₀	OH	H	OH	OH	CH ₃	"
	6. Deoxypneumocandin B ₂	OH	H	H	H	CH ₃	"
	7. Deoxypneumocandin C ₀	OH	H	OH	OH	CH ₃	"
15	8. Deoxymulundocandin	OH	H	OH	OH	H	12-Methyl tetra- decanoyl

which consists of a single step selective reduction of C4-htyr (homotyrosine) hydroxyl group of echinocandins to their monodeoxy analogues under neutral conditions without prior protection / deprotection of the equally facile C5-Orn (ornithine) hydroxyl group and purification of the monodeoxy compound from the crude reaction mixture.

2. A process as claimed in claim 1, wherein Mulundocandin is converted to Deoxymulundocandin.

3. A process as claimed in claims 1 or 2, wherein the reduction reaction is carried out by hydrogenolysis with Raney nickel in ethanol at pH 7 and room temperature.

4. A process as claimed in claims 1 to 3, wherein the hydrogenolysis is carried out in the ratio of 6.8 ml of Raney nickel per millimole of mulundocandin.




Figure 1




Figure 2

Abstract

A process for the conversion of echinocandin class of peptides to their C4 homotyrosine monodeoxy analogues

5

The invention relates to a process for the conversion of echinocandin class of peptides to their C4-homotyrosine monodeoxy analogues, particularly mulundocandin to deoxy-mulundocandin, which consists of a single step selective reduction of C4-htyr (homotyrosine) hydroxyl group of echinocandins to their monodeoxy analogues under neutral conditions without prior protection /
10 deprotection of the equally facile C5-Orn (ornithine) hydroxyl group and purification of the monodeoxy compound from the crude reaction mixture.

15